

The calculation of a normal force between multiparticle contacts using fractional operators

J.S. Leszczynski

*Institute of Mathematics & Computer Science, Technical University of
Czestochowa, 42-200 Czestochowa, ul. Dabrowskiego 73, Poland*

Keywords: granular material; molecular dynamics; discrete element model; multiparticle contacts; fractional derivative; Riemann-Liouville derivative; Caputo derivative

Abstract

This paper deals with the complex problem of how to simulate multiparticle contacts. The collision process is responsible for the transfer and dissipation of energy in granular media. A novel model of the interaction force between particles has been proposed and tested. Such model allows us to simulate multiparticle collisions and granular cohesion dynamics.

1. Introduction

Behaviours of granular materials have generated much interest in a lot of industrial processes involving transport of particles and also in natural phenomena. The key aspect in such media is how to model the interactions that may eventually take place between the particles. The collision processes are responsible for the transfer and dissipation of energy in granular materials. Moreover, the understanding of interaction process is important in order to develop simulations and theoretical

studies. Discrete models much better simulate the collision process than continuum models. We will focus on the molecular dynamics models in which takes into account an expression for the repulsive force acting between the particles. Particularly, we will analyse what happens with the multiparticle contacts. Multiparticle interactions occur when a particle contacts with surrounding particles. Typical molecular dynamics models [1, 3] are valid for particle collisions being independent from one another. Nevertheless, we observe opposite situation in behaviour of dense granular media especially for cohesive particles. In literature [6] notices the lack of energy dissipation in molecular dynamics models. We will propose novel form of a force between contacting particles and we will investigate its properties.

2. Modelling multiparticle interactions

We consider a set of spherical particles moving under optional extortion. We neglect a particle rotation. Figure 1a shows situation when particles move without any interactions. We describe a distinguished particle through its characteristic radius r_i , mass m_i , position $\mathbf{x}_i$ and linear speed $\dot{\mathbf{x}}_i$. Index i denotes a given particle and varies from 1 to np , where np is a total number of particles. Typical equation of motion for one particle without collision can be written as

$$m_i \ddot{\mathbf{x}}_i = \sum_{l=1}^{nc} \mathbf{F}_l, \quad (1)$$

where $\mathbf{F}_l$ is an optional force and nc notices a total number of the optional forces. We can distinguish the force as: gravitational one, drag force, etc. Figure 1b presents another situation, this means multiparticle collisions. We add to Eq (1)

a sum of collisional forces and we have

$$m_i \ddot{\mathbf{x}}_i + \sum_{j=1 \wedge j \neq i}^{nd_i} \mathbf{P}_j = \sum_{l=1}^{nc} \mathbf{F}_l, \quad (2)$$

where $\mathbf{P}_j$ is a collisional force acting between two particles. Temporary index j denotes a particle contacting with the given particle i and nd_i is a total number of particles surrounding and contacting with the particle i . In the molecular dynamics models, particles virtually overlap when a contact occurs. Let C_j be a point in which two particles contact as shown on the detail A in Figure 1b. Let Π be a plane tangent to colliding particles at the point C_j and ζ_j be a normal direction to Π . We decompose the collisional force into $\mathbf{P}_j = \mathbf{P}_{\xi j} + \mathbf{P}_{\eta j} + \mathbf{P}_{\zeta j}$, where $\mathbf{P}_{\zeta j}$ is the force acting on the normal direction and $\mathbf{P}_{\xi j}$, $\mathbf{P}_{\eta j}$ are forces acting on the tangent direction. In our considerations we neglect the tangent forces assuming $\mathbf{P}_{\xi j} = \mathbf{P}_{\eta j} = 0$. Here we present some examples of the normal force models which are frequently applied ones in practical simulations. Cundall and Strack [1] proposed the force being a linear combination of viscous and elastic terms

$$\mathbf{P}_{\zeta j} = c_j \cdot \dot{\zeta}_j + k_j \cdot \zeta_j, \quad (3)$$

where k_j is a stiffness of a spring whose elongation is ζ_j and c_j is a damping constant. According to the detail A presented in Figure 1b we define the virtual overlap as

$$\zeta_j = (r_j + r_i) \cdot \mathbf{e}_{\zeta j} - (\mathbf{x}_j - \mathbf{x}_i), \quad (4)$$

and a unitary vector $\mathbf{e}_{\zeta j}$ normal to Π

$$\mathbf{e}_{\zeta j} = \frac{\mathbf{x}_j - \mathbf{x}_i}{\|\mathbf{x}_j - \mathbf{x}_i\|}. \quad (5)$$

Lee [3] investigated a nonlinear version of Eq (3)

$$\mathbf{P}_{\zeta j} = c_j \cdot \dot{\zeta}_j + k_j \cdot |\zeta_j|^{\frac{3}{2}} \cdot \text{sign}(\zeta_j). \quad (6)$$

As a remark, we want to comment that Eqs (3) and (6) are typical for binary collisions. Several coefficients c_j and k_j assumed as a function of normal restitution coefficient. In the case of multiparticle contacts we have to assume the same collisional time which is not suitable for granular cohesion dynamics. Independently on physical properties of granular materials, we cannot change surface properties of contacting particles in above models. Therefore, we cannot perform simulations taking into account the particles cohesion. The restitution coefficient informs us about some work of deformation between contacting bodies but does not inform how much time is needed during particle collisions. In real behaviour of moving particles we can easy change surface properties of granular materials. Especially, when we consider the interactions in a granular material for dry particles and for wet ones. In crucial point of our discussion, we assume that momentum and energy transfer between multiparticle contacts is identified by *memory effect*. From the other hand, a given particle have to remember about surrounding particles during collision process. Using fractional calculus [5] and the generalised viscoelastic model [7] we propose a novel form of the normal force

$$\mathbf{P}_{\zeta j} = c_j^{\alpha_j} \cdot k_j^{1-\alpha_j} \cdot {}_{t_j^*}D_t^{\alpha_j}(\zeta_j), \quad (7)$$

where c_j and k_j have the same meaning like in previous models, α_j is a real order of differentiation which belongs to the range $\alpha_j \in \langle 0 \dots 1 \rangle$ and ${}_{t_j^*}D_t^{\alpha_j}(\zeta_j)$ is

a differential operator of the fractional order α_j . According to [5] we introduce a definition of such operator as left side Riemann-Liouville fractional derivative

$${}_{t_j^*}D_t^{\alpha_j}(\zeta_j) = \begin{cases} \frac{1}{\Gamma(n-\alpha_j)} \frac{d^n}{dt^n} \int_{t_j^*}^t \frac{\zeta_j(\tau)}{(t-\tau)^{\alpha_j+1-n}} d\tau & \text{for } n-1 < \alpha_j < n \\ \frac{d^n}{d(t-t_j^*)^n} \zeta_j(t) & \text{for } \alpha_j = n \end{cases}, \quad (8)$$

where $n = [\alpha_j] + 1$ and $[\cdot]$ denotes integer part of a real number. We also have a symbol t_j^* which determines begin of the collision process. Taking into consideration a fact, given by Hilfer [2], that the Riemann-Liouville derivative has no physical interpretation - especially when we try to introduce initial conditions - we formulate another one fractional derivative

$${}_t^C D_{t_j^*}^{\alpha_j}(\zeta_j) = \begin{cases} \frac{1}{\Gamma(n-\alpha_j)} \int_{t_j^*}^t \frac{\frac{d^n \zeta_j(\tau)}{d\tau^n}}{(t-\tau)^{\alpha_j+1-n}} d\tau & \text{for } n-1 < \alpha_j < n \\ \frac{d^n}{d(t-t_j^*)^n} \zeta_j(t) & \text{for } \alpha_j = n \end{cases}, \quad (9)$$

which is called Caputo derivative [2, 5]. According to [5] we describe transition between Eq (8) and Eq (9) in form

$${}_{t_j^*}D_t^{\alpha_j}(\zeta_j) = \sum_{l=0}^{n-1} \frac{(t-t_j^*)^{l-\alpha_j}}{\Gamma(l-\alpha_j+1)} \cdot \zeta_j(t_j^*) + {}_t^C D_{t_j^*}^{\alpha_j}(\zeta_j), \quad (10)$$

where the sum means initial conditions. We explain with details a meaning and dependence between these two derivatives because many investigators have concentrated on the Riemann-Liouville derivative, but this is a lack in physical applications. Everywhere in the contact process, we numerically solve Eq (2) with formula (7) together with Eq (10) by application the decomposition method described in [4].

3. Computational results

To illustrate profits of the force model given by formula (7) we compare its behaviour with the force models presented by Eqs (3) and (6). In this case we simulate a particle vertically falling down to a bottom plate as shown in Figure 2. When performing this simulation, the particle falls under gravity (we used Eq (1)) and the contact occurs in the plate (we used Eq (2)). Figure 2 presents vertical displacement of the particle over time. When the collision occurs, we apply different schemes of the normal forces: continuous line represents the linear force (3), dashed line shows the nonlinear one (6) and two dash-dot lines are responsible for our formula (7). For determined physical properties of the particle and initial conditions we observe very good agreement between our force model and the linear one. In such case we establish the parameter α in Eq (7) $\alpha = 0.051$. In the next case we can see quite good agreement between our force model and the nonlinear one when $\alpha = 0.19$. We can simulate both cases changing the non-integer order α in the fractional operator. In analysis of such behaviour we can observe that the parameter α models the surface properties. When $\alpha = 0$ we have an elastic collision but $\alpha = 1$ determines a viscous collision. We notice that the material parameters like c_j and k_j in formula (7) are constant. Therefore, the fractional order α is a surface parameter balancing between elastic and viscous properties of two contacting bodies. Moreover, formula (7) is suitable for multiparticle contacts. Figure 3 presents some behaviour of three particles in two dimensional space when the parameter α changed from 0.01 to 0.9. Such simulation is so far from reality because we neglect gravity force, tangential forces under particle contacts and particle rotations. Thin lines represents particle trajectories but thick

lines are common trajectories when particles move like a one body. In the low values of α we do not observe common trajectories. When the parameter α increases up to $\alpha = 0.9$ we can see common trajectories of two or three particles in dependence on mutual positions. We strongly notice that formula (7) is more flexible to model cohesion processes than other ones. Extending our considerations we observe different lengths of the common trajectories. This fact issues from different begins of contacting times t_j^* between two interacting particles. Therefore we can simulate multiparticle contacts in which we do not assume the same collisional time.

4. Conclusions

We have proposed and discussed a novel model of the normal force used in simulations of the particle collisions. With this model, it is possible to simulate multiparticle contacts in which we do not assume the same collisional time. This feature, in comparison to the linear and nonlinear force models is advantage of some generalisation between particle interactions. However, some of the parameters of this model may still be tuned, as for example, the parameter α in order to reflect real surface properties of contacting particles.

References

- [1] Cundall PA, Strack ODL. A discrete numerical model for granular assemblies. *Geotechnique* 1979; 29: 47-65.
- [2] Hilfer R. *Applications of fractional calculus in physics*. Singapore: World Scientific, 2000.

- [3] Lee J. Density waves in the flows of granular media. *Physical Review E* 1994; 49: 281-298.
- [4] Leszczynski JS, Ciesielski M. A numerical method for solution of ordinary differential equations of fractional order. *Lecture Notes in Computer Science* 2001; 2328: 695-702.
- [5] Oldham KB, Spanier J. The fractional calculus. Theory and applications of differentiation and integration to arbitrary order. New York: Academic Press, 1974.
- [6] Pournin L, Liebling ThM. Molecular dynamics force models for better control of energy dissipation in numerical simulations of dense granular media. *Physical Review E* 2001; 65: 011302-1-011302-7.
- [7] Schiessel H, Metzler R, Blumen A and Nonnenmacher TF. Generalized viscoelastic models: their fractional equations with solutions. *Journal of Physics A: Math. Gen.* 1995; 28: 6567-6584.




Figure 1

Sketch to illustrate particles behaviour: a) without collision; b) with multiparticle contacts.




Figure 2

Displacement over time of a particle falling down on a bottom plate.




Figure 3

Multiparticle contacts in dependence on surface properties.